

Original link: [http://www.uptodate.com/contents/high-fiber-diet-beyond-the-basics?source=search\\_result&search=high+fiber+diet&selectedTitle=1~150](http://www.uptodate.com/contents/high-fiber-diet-beyond-the-basics?source=search_result&search=high+fiber+diet&selectedTitle=1~150)

## Patient information: High-fiber diet (Beyond the Basics)

### HIGH-FIBER DIET OVERVIEW

Eating a diet that is high in fiber has many potential health benefits, including a decreased risk of heart disease, stroke, and type 2 diabetes. Because high-fiber foods may be healthy for reasons other than their fiber content, the research has not always been able to determine if fiber is the healthful component. A high-fiber diet is a commonly recommended treatment for digestive problems, such as constipation, diarrhea, and hemorrhoids, although individual results vary widely, and the scientific evidence supporting these recommendations is weak.

Fiber is normally found in beans, grains, vegetables, and fruits. However, most people do not eat as much fiber as is commonly recommended. This topic discusses what fiber is, why it is helpful, and how to increase dietary fiber.

### WHAT IS FIBER?

There is no single dietary "fiber." Traditionally, fiber was considered that substance found in the outer layers of grains or plants and which was not digested in the intestines. Wheat bran, the outer layer of wheat grain, fit this model. We now know that "fiber" actually consists of a number of different substances. The term "dietary fiber" includes all of these substances and is now considered a better term than just "fiber."

Most dietary fiber is not digested or absorbed, so it stays within the intestine where it modulates digestion of other foods and affects the consistency of stool. There are two types of fiber, each of which is thought to have its own benefits:

- Soluble fiber consists of a group of substances that is made of carbohydrates and dissolves in water. Examples of foods that contain soluble fiber include fruits, oats, barley, and legumes (peas and beans).
- Insoluble fiber comes from plant cells walls and does not dissolve in water. Examples of foods that contain insoluble fiber include wheat, rye, and other grains. The traditional fiber, wheat bran, is a type of insoluble fiber.
- Dietary fiber is the sum of all soluble and insoluble fiber.

### BENEFITS OF A HIGH-FIBER DIET

The health effects of a high-fiber may depend to some extent on the type of fiber eaten. However, the difference between the health effects of two types of fiber are not very clear and

may vary between individuals, so many providers encourage adding fiber in whatever way is easiest for the patient.

There are several potential benefits of eating a diet with high-fiber content:

- Insoluble fiber (wheat bran, and some fruits and vegetables) has been recommended to treat digestive problems such as constipation, hemorrhoids, chronic diarrhea, and fecal incontinence. Fiber bulks the stool, making it softer and easier to pass. Fiber helps the stool pass regularly, although it is not a laxative. (See ["Patient information: Constipation in adults \(Beyond the Basics\)"](#) and ["Patient information: Hemorrhoids \(Beyond the Basics\)"](#) and ["Patient information: Chronic diarrhea in adults \(Beyond the Basics\)"](#).)
- Soluble fiber (psyllium, pectin, wheat dextrin, and oat products) can reduce the risk of coronary artery disease and stroke by 40 to 50 percent (compared to a low fiber diet) [1,2].
- Soluble fiber can also reduce the risk of developing type 2 diabetes. In people who have diabetes (type 1 and 2), soluble fiber can help to control blood glucose levels.
- It is not clear if a high-fiber diet is beneficial for people with irritable bowel syndrome or diverticulosis. Fiber may be helpful for some people with these diagnoses while it may worsen symptoms in others.

## HOW MUCH FIBER DO I NEED?

The recommended amount of dietary fiber is 20 to 35 grams per day. By reading the nutrition label on packaged foods, it is possible to determine the number of grams of dietary fiber per serving ([figure 1](#)).

Dietary sources of fiber — The fiber content of many foods, including fruits and vegetables, is available in the table ([table 1](#)). Breakfast cereals can be a good source of fiber. Some fruits and vegetables are particularly helpful in treating constipation, such as prunes and prune juice.

Other sources of fiber — For those who do not like high-fiber foods such as fruits, beans, and vegetables, a good source of fiber is unprocessed wheat bran; one to two tablespoons can be mixed with food. One tablespoon of wheat bran contains approximately 1.6 grams of fiber.

In addition, a number of fiber supplements are available. Examples include psyllium, methylcellulose, wheat dextrin, and calcium polycarbophil. The dose of the fiber supplement should be increased slowly to prevent gas and cramping, and the supplement should be taken with adequate fluid. The fiber in these supplements is mostly of the soluble type.

## FIBER SIDE EFFECTS

Adding fiber to the diet can have some side effects, such as abdominal bloating or gas. This can sometimes be minimized by starting with a small amount and slowly increasing until stools become softer and more frequent.

However, many people, including those with irritable bowel syndrome, cannot tolerate fiber supplements and do better by not increasing fiber in their diet. (See ["Patient information: Irritable bowel syndrome \(Beyond the Basics\)"](#).)

## WHERE TO GET MORE INFORMATION

Your healthcare provider is the best source of information for questions and concerns related to your medical problem.

This article will be updated as needed on our web site ([www.uptodate.com/patients](http://www.uptodate.com/patients)). Related topics for patients, as well as selected articles written for healthcare professionals, are also available. Some of the most relevant are listed below.

Patient level information — UpToDate offers two types of patient education materials.

The Basics — The Basics patient education pieces answer the four or five key questions a patient might have about a given condition. These articles are best for patients who want a general overview and who prefer short, easy-to-read materials.

[Patient information: High-fiber diet \(The Basics\)](#)

[Patient information: Managing loss of appetite and weight loss with cancer \(The Basics\)](#)

[Patient information: Diet and health \(The Basics\)](#)

[Patient information: Fecal incontinence \(The Basics\)](#)

[Patient information: Diabetes and diet \(The Basics\)](#)

[Patient information: Rectal prolapse in adults \(The Basics\)](#)

[Patient information: Colostomy care \(The Basics\)](#)

Beyond the Basics — Beyond the Basics patient education pieces are longer, more sophisticated, and more detailed. These articles are best for patients who want in-depth information and are comfortable with some medical jargon.

[Patient information: Constipation in adults \(Beyond the Basics\)](#)

[Patient information: Hemorrhoids \(Beyond the Basics\)](#)

[Patient information: Chronic diarrhea in adults \(Beyond the Basics\)](#)

[Patient information: Irritable bowel syndrome \(Beyond the Basics\)](#)

Professional level information — Professional level articles are designed to keep doctors and other health professionals up-to-date on the latest medical findings. These articles are thorough, long, and complex, and they contain multiple references to the research on which they are based. Professional level articles are best for people who are comfortable with a lot of medical terminology and who want to read the same materials their doctors are reading.

[Colorectal cancer: Epidemiology, risk factors, and protective factors](#)

[Diet in the treatment and prevention of hypertension](#)

[Lipid lowering with diet or dietary supplements](#)

[Healthy diet in adults](#)

[Management of chronic constipation in adults](#)

The following organizations also provide reliable health information.

- National Library of Medicine

([www.nlm.nih.gov/medlineplus/dietaryfiber.html](http://www.nlm.nih.gov/medlineplus/dietaryfiber.html), available in Spanish)

- National Institute on Diabetes and Digestive and Kidney Diseases

([www.niddk.nih.gov](http://www.niddk.nih.gov))

- Harvard School of Public Health

([www.hsph.harvard.edu/nutritionsource/what-should-you-eat/fiber/index.html](http://www.hsph.harvard.edu/nutritionsource/what-should-you-eat/fiber/index.html))

[1-5]

Literature review current through: Oct 2013. | This topic last updated: Jul 19, 2013.

[Find Print](#)

The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of this website is governed by the [UpToDate Terms of Use](#) ©2013 UpToDate, Inc.

## References

▲[Top](#)

1. [Gillman MW, Cupples LA, Gagnon D, et al. Protective effect of fruits and vegetables on development of stroke in men. JAMA 1995; 273:1113.](#)
2. [Jensen MK, Koh-Banerjee P, Hu FB, et al. Intakes of whole grains, bran, and germ and the risk of coronary heart disease in men. Am J Clin Nutr 2004; 80:1492.](#)
3. [Negri E, Franceschi S, Parpinel M, La Vecchia C. Fiber intake and risk of colorectal cancer. Cancer Epidemiol Biomarkers Prev 1998; 7:667.](#)
4. [Willett WC. Diet and cancer: an evolving picture. JAMA 2005; 293:233.](#)
5. [Suarez NC, Ford AC. Systematic review: the effects of fibre in the management of chronic idiopathic constipation. Aliment Pharmacol Ther 2011; 33:895.](#)